

DISTURBED SEXUAL CHARACTERISTICS IN MALE MOSQUITOFISH (*GAMBUSIA HOLBROOKI*) FROM A LAKE CONTAMINATED WITH ENDOCRINE DISRUPTERS

Gunnar Toft, Thea M. Edwards, Erik Baatrup, and Louis J.
Guillette Jr.

doi:10.1289/ehp.6022 (available at <http://dx.doi.org/>)

Online 23 January 2003



**DISTURBED SEXUAL CHARACTERISTICS IN MALE MOSQUITOFISH (*GAMBUSIA
HOLBROOKI*) FROM A LAKE CONTAMINATED WITH ENDOCRINE DISRUPTERS**

Gunnar Toft ^{1,2*}, Thea M. Edwards ², Erik Baatrup ¹ and Louis J. Guillette Jr. ²

¹Institute of Biological Sciences

University of Aarhus

Building 135

DK-8000 Aarhus C

Denmark

²Department of Zoology

223 Bartram Hall

University of Florida

Gainesville, FL, 32611-8525

USA.

*Current address of corresponding author: Gunnar Toft, Department of Occupational Medicine,
Aarhus University Hospital, Noerrebrogade 44, build. 2C, DK-8000 Aarhus C. Denmark.
Phone: 45 8949 4251; Fax: +45 8949 4260; e-mail: gutof@akh.aaa.dk.

Running title

Altered sexual characteristics in mosquitofish.

Key words

Antiandrogen, courtship behavior, endocrine disruption, estradiol, mosquitofish, sex characters, sperm count, testosterone.

Acknowledgments

This work was supported by the Danish Natural Science Research Council and the Danish Environmental Research Programme. This work was also supported, in part, by grant # CR826357-01-1 from the US EPA to LJG.

List of Abbreviations

DDD: 1,1-dichloro-2,2-bis(*p*-chlorophenyl)ethane

DDE: 1,1-dichloro-2,2-bis(*p*-chlorophenyl)ethylene

DDT: 1,1,1-trichloro-2,2-bis(*p*-chlorophenyl)ethane

EDC : Endocrine disrupting compound

RIA: Radioimmunoassay

MS-222: 3-Aminobenzoic acid ethyl ester

Outline of manuscript section headers

Abstract

Introduction

Materials and Methods

Results

Discussion

References

Abstract

Previous laboratory studies have demonstrated that estrogenic and antiandrogenic chemicals can alter several sexual characteristics in male poeciliid fishes. Whether similar disturbances occur under field conditions remains to be confirmed. Lake Apopka (Florida, USA) is contaminated with numerous chemicals, some of which possess endocrine disrupting activity. Male mosquitofish (*Gambusia holbrooki*) were collected monthly from December 2000 through May 2001 from Lake Apopka and two nearby reference lakes, Orange Lake and Lake Woodruff National Wildlife Refuge. Selected sexual characteristics were compared temporally and among lakes during the collection period. Male fish from Lake Apopka had slightly shorter gonopodia and on average 32% and 47% fewer sperm cells per mg testis, when compared to the fish collected from Orange Lake and Lake Woodruff, respectively. The testes weights increased markedly during spring with significantly smaller testes in fish from Lake Apopka than from Orange Lake, but surprisingly, the smallest testes occurred in males obtained from the Lake Woodruff population. The highest liver weights were found in the Lake Apopka population. Whole body concentrations of testosterone and estradiol varied among months; peak testosterone concentration occurred in January and was significantly lower in male fish from Lake Apopka compared to Orange Lake. The intensity of male courtship behavior was highly correlated to body testosterone concentration, but no statistically significant differences in sexual behavior among the lakes were found. It is concluded that sexual characteristics of relevance to male reproductive capacity are altered in the Lake Apopka mosquitofish population. The presence of chemicals with antiandrogenic effects in Lake Apopka as a possible cause of the observed alterations is discussed.